## If a person is experiencing COVID-19 symptoms\*, get the

# **DIAGNOSTIC TEST**



### HOW TO GET TESTED

Discuss symptoms with your health care provider and get a lab order for testing

# HOW THE TEST IS COVERED THROUGH INDEPENDENCE

Covered at authorized labs with no costsharing under all commercial medical plans (includes fully-insured and self-funded plans)

# USEFUL FOR EMPLOYERS IN WORKPLACE REOPENING?

Not helpful in most workplaces. If employees are sick they should stay home. Employees could be temperature screened along with other space planning precautions.

#### MOLECULAR DIAGNOSTIC TEST

Also known as a diagnostic test, viral test, nucleic acid amplification tests (NAAT), or RT-PCR test



Indicates active infection



Current molecular tests are more accurate than antigen tests





Results take 1-7 days



#### ANTIGEN DIAGNOSTIC TEST

Also known as a rapid diagnostic test



Detects specific proteins on the surface of the virus



More likely to miss active infection than molecular tests



Nasal or throat swah





<sup>\*</sup>Symptoms include fever, cough, shortness of breath, difficulty breathing, body aches, headache, loss of taste, sore throat, congestion

A person who suspects they had COVID-19 might consider getting the

# **ANTIBODY TEST**

Also known as the serological test, serology blood test



### HOW TO GET TESTED

Get a lab order from your health care provider

## HOW THE TEST IS COVERED THROUGH INDEPENDENCE

Covered at authorized labs with no cost-sharing under all commercial medical plans (includes fully-insured and self-funded plans)

# USEFUL FOR EMPLOYERS IN WORKPLACE REOPENING?

Not recommended, according to CDC guidelines



**Population Health Benefit:** A large group of immune individuals inhibits transmission of the virus. 'Herd immunity' is projected when about 60% of the population exhibits immunity



Looks for antibodies that are made by the immune system. Antibodies may stay in the body for several weeks after recovery. It is unknown how strong the immunity will be or how long the protection will last.



This test cannot diagnose active infection



Finger stick or blood draw



Results take 1-3 days



## CDC TIPS FOR STOPPING THE SPREAD

- Wash hands frequently
- Stay 6' apart
- Avoid large groups
- Wear face masks

For the latest COVID-19 information go to ibx.com/COVID19.

